

# CraveArrest™



Natural support for neurotransmitter production

By David Brady, ND, DC, CCN, DACBN & Suzanne Copp, MS

THIS INFORMATION IS PROVIDED FOR THE USE OF PHYSICIANS AND OTHER LICENSED HEALTH CARE PRACTITIONERS ONLY. THIS INFORMATION IS INTENDED FOR PHYSICIANS AND OTHER LICENSED HEALTH CARE PROVIDERS TO USE AS A BASIS FOR DETERMINING WHETHER OR NOT TO RECOMMEND THESE PRODUCTS TO THEIR PATIENTS. THIS MEDICAL AND SCIENTIFIC INFORMATION IS NOT FOR USE BY CONSUMERS. THE DIETARY SUPPLEMENT PRODUCTS OFFERED BY DESIGNS FOR HEALTH ARE NOT INTENDED FOR USE BY CONSUMERS AS A MEANS TO CURE, TREAT, PREVENT, DIAGNOSE, OR MITIGATE ANY DISEASE OR OTHER MEDICAL CONDITION.

Many factors have an impact on brain neurochemistry and endocrine balance such as age, genetics, circadian rhythms and various environmental factors including stress, diet, exercise and medications. Many people do not synthesize adequate serotonin including diabetics and those on SSRIs for an extended period of time. Dieting is known to lower levels of tryptophan, an amino acid needed to make serotonin. Dieters, as well as others, often need help in controlling food cravings, mood swings and addictions. Cravings for cigarettes are known to be controlled by dopamine receptors. Dopamine is another very important neurotransmitter often low in those who are prone to addictive or erratic behavior. Imagine what might be thrown off if one or more of these important neurotransmitters is deficient.

## The perception of hunger is determined by many physiologic factors:

1. Neurotransmitter balance of serotonin, dopamine, GABA, glutamine, acetylcholine and opioids. In general, any deficiency of these could trigger anxiety, hunger, depression, and cravings.
2. Genetic specificity of brain neurochemistry and hormonal balance.
3. Body fat stores. Adipose tissue produces various metabolic signals that affect the appetite: leptin helps to control hunger, inflammatory cytokines (TNF-alpha, IL-6, etc.) increase it.
4. The level of glycogen stores in the liver is communicated to the brain through the vagal nervous system, thus attempting to influence the appetite.
5. Blood glucose levels are perceived directly by the brain through glucoreceptors. Low blood glucose levels trigger carbohydrate cravings.
6. Adrenal hormones, cortisol and adrenaline, increase hunger and the production of sex hormones (estrogen, testosterone, progesterone). These all influence appetite.
7. Gut-derived hormones are produced directly in response to the macronutrient composition and size of the meals: cholecystekinin (CCK), in response to protein, fat and the stretching of the stomach by food/drink volume; galanin, in response to fat; ghrelin, in response to an empty stomach. They signal the brain directly. CCK and galanin reduce hunger while ghrelin stimulates it.
8. Pancreatic hormones: high insulin levels signal the brain to reduce hunger if the tissues are insulin sensitive.

The amino acid L-tryptophan (LT) converts into 5-hydroxytryptophan (5-HTP) which converts into 5-hydroxytryptamine (5-HT), also known as serotonin. 5-HTP readily enters the blood brain barrier and makes serotonin. Oral administration of 5-HTP has been shown to successfully raise serotonin levels. Serotonin is known to control sleep, depression, anxiety, aggression, appetite, temperature, sexual behavior and pain sensation.

- Other neurotransmitters and CNS chemicals such as melatonin, dopamine, norepinephrine, and beta-endorphin have all been shown to increase following oral administration of 5-HTP.
- This ability to increase not only serotonin levels in the brain, but also dopamine and norepinephrine, allows 5-HTP to produce some significant and unique effects on brain chemistry and on serotonin-related conditions which other substances, including L-tryptophan, cannot duplicate.
- L-tryptophan may raise serotonin levels or may be shunted into the synthesis of niacin or the production of protein.

Research shows that type II diabetics have lower brain tryptophan levels and higher rates of depression. This may help explain their propensity towards sugar addiction beyond the typical fluctuating blood glucose levels. When twenty overweight NIDDM patients were given 750 mg/day of 5-HTP or placebo for two weeks in a double-blind study, their daily energy intake decreased considerably as well as their carbohydrate and fat intake. Body weight also diminished along with appetite.<sup>29</sup>

## Supplement Facts

Serving Size 2 capsules

Servings Per Container 60

Amount Per Serving	% Daily Value	
Vitamin C (as Ascorbic Acid)	100 mg	170%
Niacin (Vitamin B-3)(as Niacinamide)	20 mg	100%
Vitamin B-6 (as Pyridoxine HCL 15 mg; 20 mg Pyridoxal-5-Phosphate 5 mg)	20 mg	1000%
Vitamin B-12 (as Methylcobalamin)	25 mcg	420%
L-Tyrosine	1000 mg	*
5-HTP (5-Hydroxytryptophan)	100 mg	*
Taurine	50 mg	*
Rhodiola ( <i>Rhodiola rosea</i> )(root)	50 mg	*
[standardized to contain 3% rosavins and 1% salidroside]		

\*Daily Value not established.

Other Ingredients: Microcrystalline cellulose, vegetable stearate.

## Increasing neurotransmitters may be helpful for:

- Food cravings
- Appetite control
- Smoking cessation
- Insomnia
- Depression
- Anxiety
- Mood swings
- Premenstrual syndrome
- Obsessive compulsive disorders
- Addictions
- Stress
- Type II diabetes

## MECHANISMS OF ACTION OF BRAIN NUTRIENTS

NUTRIENT	ACTION	REFERENCE
<b>Vitamin C 100 mg</b>	• Cofactor in adrenaline synthesis	<b>1, 2</b>
<b>Vitamin B6 20 mg</b>	• Cofactor in serotonin and dopamine synthesis	<b>4, 5</b>
<b>Vitamin B12 25 mcg</b>	• Cofactor in serotonin and dopamine synthesis	<b>3, 28</b>
<b>Niacinamide 20 mg</b>	• Cofactor in brain energy production	<b>7</b>
<b>L-Tyrosine 1,000 mg</b>	<ul style="list-style-type: none"> <li>• Precursor to dopamine and adrenaline</li> <li>• Antidepressant</li> <li>• Reduces cravings for tobacco</li> </ul>	<b>13, 14 15 16</b>
<b>5-Hydroxytryptophan 100 mg</b>	<ul style="list-style-type: none"> <li>• Precursor to serotonin and melatonin</li> <li>• Reduces carbohydrate cravings and appetite</li> <li>• Corrects age-related decline in serotonin</li> </ul>	<b>1, 28 22, 23 11</b>
<b>Taurine 50 mg</b>	<ul style="list-style-type: none"> <li>• Important regulator of calcium and neurotransmitters within the heart, muscles and brain</li> <li>• Helps calm the nervous system by regulating neurotransmitters</li> <li>• Strengthens and protects healthy cell membranes</li> </ul>	<b>1 8, 9, 10 12</b>
<b>Rhodiola Rosea 50 mg</b>	<ul style="list-style-type: none"> <li>• Improves levels and metabolism of:                             <ul style="list-style-type: none"> <li>- beta endorphin</li> <li>- dopamine</li> <li>- serotonin</li> </ul> </li> <li>• Adaptogen action, helping with stress</li> </ul>	<b>21 21</b>

### References

1. Barbeau, A. The Brain, The Heart and Taurine. *Can J Neurol Sci* 1975;2(4):346.
2. Lee L, Kang SA, Lee HO, Lee BH, Jung IK, Lee JE, Hoe YS. Effect of supplementation of vitamin E and vitamin C on brain acetylcholinesterase activity and neurotransmitter levels in rats treated with scopolamine, an inducer of dementia. *J Nutr Sci Vitaminol (Tokyo)* 2001 Oct;47(5):323-8.
3. Petrie WM, Ban TA. Vitamins in psychiatry. Do they have a role? *Drugs* 1985 Jul;30(1):58-65.
4. Bhagavan HN, Coleman M, Coursin DB. The effect of pyridoxine hydrochloride on blood serotonin and pyridoxal phosphate contents in hyperactive children. *Pediatrics* 1975 Mar;55(3):437-41.
5. Bernstein AL. Vitamin B6 in clinical neurology. *Ann N Y Acad Sci* 1990;585:250-60.
6. Kan JB, Chouvet G, Hery F, Debilly G, Mermet A, Glowinski J, Pujol JF. Daily variations of various parameters of serotonin metabolism in the rat brain. Circadian variations of tryptophan-5-hydroxylase in the raphe nuclei and the striatum. *Brain Res* 1977 Mar 4;123(1):125-36.
7. Scherer B, Kramer W. Influence of niacinamide administration on brain 5-HT and a possible mode of action. *Life Sci* 1972 Feb 15;11(4):189-95.
8. Ashinara, M. Immunohistochemical Study of The Human Olfactory System. *Nippon Jibiinkoka Gakkai Kaiho* 1992;985(6):851-9.
9. Gaull, B.E. and Rassin D.K. Taurine in Development and Nutrition. *Sulphur in Biology Ciba Foundation Symposium*, 1980.
10. Hayes, K.C. Taurine Requirement in Primates. *Nutr Rev* 1985;43(3):65-70.
11. Neurochemical changes related to ageing in the rat brain and the effect of DL-alpha-lipoic acid. Arivazhagan P, Panneerselvam C. *Exp Gerontol* 2002 Dec;37(12):1489-94.
12. Chapman, G.W. and Greenwood, C.E. Taurine in Nutrition and Brain Development. *Nutri Res* 1988;8:955-68.
13. Lehnert H, Wurtman RJ. Amino acid control of neurotransmitter synthesis and release: physiological and clinical implications. *Psychother Psychosom* 1993;60(1):18-32.
14. Conlay LA, Zeisel SH. Neurotransmitter precursors and brain function. *Neurosurgery* 1982 Apr;10(4):524-9.
15. Fernstrom JD. Dietary precursors and brain neurotransmitter formation. *Annu Rev Med* 1981;32:413-25.
16. McCarty MF. Nutritional support of central catecholaminergic tone may aid smoking withdrawal. *Med Hypotheses* 1982 Jan;8(1):95-102.
17. McCarty MF. Promotion of hepatic lipid oxidation and gluconeogenesis as a strategy for appetite control. *Med Hypotheses* 1994 Apr;42(4):215-25.
18. Davidson JR, Abraham K, Connor KM, McLeod MN. Effectiveness of chromium in atypical depression: a placebo-controlled trial. *Biol Psychiatry* 2003 Feb 1;53(3):261-4.
19. McLeod MN, Golden RN. Chromium treatment of depression. *Int J Neuropsychopharmacol* 2000 Dec;3(4):311-314.
20. Effect of vanadium on insulin sensitivity and appetite. 20. Wang J, Yuen VG, McNeill JH. *Metabolism* 2001 Jun;50(6):667-73.
21. Kelly GS. Rhodiola rosea: a possible plant adaptogen. *Altern Med Rev* 2001 Jun;6(3):293-302.
22. Cangiano C, Ceci F, Cascino A, Del Ben M. Eating behavior and adherence to dietary prescriptions in obese adult subjects treated with 5-hydroxytryptophan. *Am J Clin Nutr* 1992 Nov;56(5):863-7.
23. Ceci F, Cangiano C, Cairella M, Cascino A, Del Ben M, Muscaritoli M, Sibilia L, Rossi Fanelli E. The effects of oral 5-hydroxytryptophan administration on feeding behavior in obese adult female subjects. *J Neural Transm* 1989;76(2):109-17.
24. Miller AL. The methionine-homocysteine cycle and its effects on cognitive diseases. 1: *Altern Med Rev* 2003 Feb;8(1):7-19.
25. Alpert JE, Fava M. Nutrition and depression: the role of folate. *Nutr Rev* 1997 May;55(5):145-9.
26. Young SN. The use of diet and dietary components in the study of factors controlling affect in humans: a review. 1: *J Psychiatry Neurosci* 1993 Nov;18(5):235-44.
27. Young SN, Ghadirian AM. Folic acid and psychopathology. *Prog Neuropsychopharmacol Biol Psychiatry* 1989;13(6):841-63.
28. Birdsall TC. 5-Hydroxytryptophan: a clinically-effective serotonin precursor. *Altern Med Rev* 1998 Aug;3(4):271-80.
29. *Int J Obes Relat Metab Disor* 1998 July; 22(7): 648-54. Effects of oral 5-hydroxytryptophan on energy intake and macronutrient selection in non-insulin dependent diabetic patients.

**To contact Designs for Health, please call us at (800) 847-8302, or visit us on the web at [www.designsforhealth.com](http://www.designsforhealth.com).**