

GI Microb-X™



Natural support for gastrointestinal health

By David M. Brady, ND, DC, CCN, DACBN & Suzanne Copp, MS

THIS INFORMATION IS PROVIDED FOR THE USE OF PHYSICIANS AND OTHER LICENSED HEALTH CARE PRACTITIONERS ONLY. THIS INFORMATION IS INTENDED FOR PHYSICIANS AND OTHER LICENSED HEALTH CARE PROVIDERS TO USE AS A BASIS FOR DETERMINING WHETHER OR NOT TO RECOMMEND THESE PRODUCTS TO THEIR PATIENTS. THIS MEDICAL AND SCIENTIFIC INFORMATION IS NOT FOR USE BY CONSUMERS. THE DIETARY SUPPLEMENT PRODUCTS OFFERED BY DESIGNS FOR HEALTH ARE NOT INTENDED FOR USE BY CONSUMERS AS A MEANS TO CURE, TREAT, PREVENT, DIAGNOSE, OR MITIGATE ANY DISEASE OR OTHER MEDICAL CONDITION.

GI Microb-X™ is a blend of botanical extracts with a long history of use as natural antimicrobials. The ingredients provide a broad spectrum of activity against the most common pathogens present in the human GI tract in conditions of dysbiosis, while being relatively sparing of normal flora. The botanical substances in GI Microb-X™ are also commonly represented on sensitivity testing provided by the major functional laboratories performing stool analysis.

GI Microb-X™ Features:

- **Tribulus Extract** (also known as puncture vine), a member of the Zygophyllaceae family, is an annual herb found in many tropical and moderate areas of the world, including the US and Mexico, the Mediterranean region, and throughout Asia. Its health and medicinal effects are the result of active phytochemicals steroidal saponins currently referred to by medical researchers and physicians as X steroidal saponins. These X steroidal saponins have the ability to influence the entire immune system of the body and have been shown to have antibacterial and antiviral effects.
- **Berberine** is a bitter-tasting, yellow, plant alkaloid with a long history of medicinal use in Chinese and Ayurvedic medicine. Berberine is present in the roots, rhizomes and stem bark of various plants including *Berberis aquifolium* (Oregon grape), *Berberis vulgaris* (barberry), *Hydrastis canadensis* (goldenseal), *Coptis chinensis* (coptis or goldenthrad), and *Berberis aristata* (tree turmeric). Berberine has also been used historically as a dye, due to its yellow color. Varma first documented the use of berberine in 1933 for the eye infection chronic trachoma. There is clinical evidence to support berberine's use in the treatment of bacterial diarrhea. Berberine has also shown antimicrobial activity against bacteria, viruses, fungi, protozoans, helminths (worms), and chlamydia.
- **Artemisinin** (from Sweet Wormwood; *Artemisia annua*) demonstrates significant antimicrobial effects and has been traditionally used even in the treatment of malaria. It has a particular historic use of being used to treat parasitic gastrointestinal infections. It is derived from the inner bark (wood) of the Artemesia annua tree and it kills parasites (worms), therefore it became known by the common name of wormwood.
- **Black Walnut Extract** (*Juglans nigra*) also has a long history of use as an intestinal anti-parasitic (i.e., vermifuge, antihelmitic) in botanical medicine. It also possesses activity against common bacterial and fungal pathogens that occur in GI dysbiosis.
- **Bearberry Extract** (*Arctostaphylos uva-ursi*) is grown throughout Asia, North America, and Europe and has a long history of medicinal use dating back to the 13th Century. The leaves have been used worldwide as a diuretic, astringent, antiseptic and treatment for urinary tract and gastrointestinal infections. A tea brewed with the leaves has also been used as a laxative. Arbutin, the main chemical constituent of uva ursi, is a phenolic glycoside that becomes hydrolyzed to hydroquinone. Both chemicals contribute to the antiseptic effects in the urinary and GI tracts. Other active constituents include tannins, mono and triterpenes, and flavonoids. Arbutin alone has also been reported to relieve pain from kidney stones, cystitis and nephritis.

Available in 60 & 120 count capsules

Supplement Facts	
Serving Size 1 capsule	
Amount Per Serving	% Daily Value
Tribulus Extract (<i>Tribulus terrestris</i>)(aerial) [standardized to contain 40% saponins]	200 mg *
Magnesium Caprylate (yielding 120 mg caprylic acid; 10 mg magnesium)	150 mg *
Berberine Sulfate (<i>Berberis aristata</i>)(root)	100 mg *
Bearberry Extract (<i>Arctostaphylos uva-ursi</i>)(leaf) [standardized to contain 20% arbutin]	100 mg *
Black Walnut Powder (<i>Juglans nigra</i>)(hull)	100 mg *
Barberry Extract (<i>Berberis vulgaris</i>)(bark) [standardized to contain 6% berberine]	50 mg *
Artemisinin (from Sweet Wormwood) (<i>Artemisia annua</i>)(herb)	15 mg *

*Daily Value not established.

Other Ingredients: Cellulose (capsule), vegetable stearate, silicon dioxide.

Contains tree nuts (walnuts).



- **Caprylic Acid** is the common name for the eight-carbon straight chain fatty acid known by the systematic name octanoic acid. It is found naturally in palm and coconut oil, and in breast milk. Health care practitioners often recommend caprylic acid for use with candidiasis and bacterial infections. Due to its relatively short chain length it has no difficulty in penetrating fatty cell wall membranes, hence its effectiveness in combating certain lipid-coated bacteria, such as *Staphylococcus aureus*, various species of *Streptococcus*, and intramucosal *Candida*. Caprylic acid may affect the fluidity of viral and fungal cell membranes. The lactoperoxidase system in combination with caprylic acid can inhibit the growth of *Escherichia coli* and *Staphylococcus aureus* in food. Studies have reported that dietary caprylic acid inhibits the growth of *Candida albicans* and other opportunistic fungi in both the small and the large intestine. At the same time, caprylic acid does not seem to adversely affect the growth of beneficial intestinal microflora.

Recommended Use

- As a dietary supplement, take one capsule per day on an empty stomach, or as directed by a health care practitioner.

References

1. Bacom A. Incorporating Herbal Medicine Into Clinical Practice. Philadelphia: F.A. Davis; 2002.
2. Chevallier A. Encyclopedia of Herbal Medicine. London: Dorling Kindersley; 2000.
3. Fetrow C, Avila J. Professional's Handbook of Complimentary & Alternative Medicines. Springhouse, PA: Springhouse; 1999.
4. Pizzorno J, Murray M. Textbook of Natural Medicine. 2nd ed. Edinburgh: Churchill Livingstone;1999.
5. Robbers J, Speedie M, Tyler V. Pharmacognosy and Pharmacobiotechnology. Philadelphia: Lippicott Williams & Wilkins; 1996.
6. Werbach M, Murray M. Botanical Influences on Illness. Tarzana, CA: Third Line Press; 1994.
7. Berberine. Altern Med Rev 2000;5(2):175-177.
8. Varma RL. Berberine sulphate in chronic trachoma. Indian Med Gaz 1933;68:122.
9. Khosla, P. K., Neeraj, V. I., Gupta, S. K., and Satpathy, G. Berberine, a potential drug for trachoma. Rev Int Trach.Pathol Ocul Trop Subtrop Sante Publique 1992;69:147-165.
10. Mohan, M., Pant, C. R., Angra, S. K., and Mahajan, V. M. Berberine in trachoma. (A clinical trial). Indian J Ophthalmol. 1982;30(2):69-75
11. Khin, Maung U., Myo, Khin, Nyunt, Nyunt Wai, Aye, Kyaw, and Tin, U. Clinical trial of berberine in acute watery diarrhoea. Br Med J (Clin Res Ed) 12-7-1985;291(6509):1601-1605.
12. Rabbani, G. H., Butler, T., Knight, J., Sanyal, S. C., and Alam, K. Randomized controlled trial of berberine sulfate therapy for diarrhea due to enterotoxigenic *Escherichia coli* and *Vibrio cholerae*. J Infect.Dis 1987;155(5):979-984.
13. Gupte, S. Use of berberine in treatment of giardiasis. Am J Dis child 1975;129(7):866.
14. Sharma R, Joshi CK, and Goyal RK. Berberine tannate in acute diarrhoea. Indian Pediatrics 1970;7(9):496-501.
15. Kamat SA. Clinical trials with berberine hydrochloride for the control of diarrhea in acute gastroenteritis. J Assoc Physicians India 1967;15:525-529.
16. Saksena HC, Tomar VN, and Soangra MR. Efficacy of a new salt of Berberine Uni-Berberine in oriental sore. Current Medical Practice 1970;14:247-252.
17. Purohit SK, Kochar DK, Lal BB, and et al. Cultivation of *Leishmania tropica* from untreated and treated cases of oriental sore. Indian Journal of Public Health 1982;26(1):34-37.
18. Edwards-Jones V, Buck R, Shawcross SG, Dawson MM, Dunn K. The effect of essential oils on methicillin-resistant *Staphylococcus aureus* using a dressing model. Burns. 2004 Dec;30(8):772-7.
19. Siegers C, Bodinet C, Ali SS, et al. Bacterial deconjugation of arbutin by *Escherichia coli*. Phytomedicine 2003;10 Suppl 4:58-60.
20. Sugai T. [Clinical effects of arbutin in patients with chloasma]. Skin Research 1992;34:522-529.
21. McLay, J. C., Kennedy, M. J., Orouke, A. L., Elliot, R. M., and Simmonds, R. S. Inhibition of bacterial foodborne pathogens by the lactoperoxidase system in combination with monolaurin. Int J Food Microbiol. 2-25-2002;73(1):1-9.
22. Neuhauser, I. Successful treatment of intestinal moniliasis with fatty acid-resin complex. AMA.Arch Intern Med 1954;93(1):53-60.

To contact Designs for Health, please call us at (800) 847-8302, or visit us on the web at www.designsforhealth.com.